

RESEARCH

Open Access

Socioeconomic factors, attitudes and practices associated with malaria prevention in the coastal plain of Chiapas, Mexico

Merit Mora-Ruiz, R Patricia Penilla*, José G Ordóñez, Alma D López, Francisco Solis, José Luis Torres-Estrada and Américo D Rodríguez

Abstract

Background: Mexico is in the malaria pre-elimination phase; therefore, continuous assessment and understanding of the social and behavioural risk factors related to exposure to malaria are necessary to achieve the overall goal. The aim of this research was to investigate socio-economic backgrounds, attitudes and practices related with malaria in rural locations from the coastal plain of Chiapas.

Methods: In January 2012, 542 interviews were conducted to householders from 20 villages across the coastal plain of Chiapas. Questions were about housing conditions, protection from mosquito bites and general information of householders. Chi² analyses were performed to see whether there was a dependence of those reported having malaria with their house conditions and their malaria preventive practices. Results were discussed and also compared statistically against those obtained 17 years ago from the same area.

Results: Most households had 2–5 people (73.6%), 91.6% of houses had 1–3 bedrooms. The physical structure of the houses consisted of walls mainly made of block or brick 72.3%, the floor made of cement 90.0%, while the roof made of zinc sheet 43.9%, and straw or palm 42.2%. A 23.1% of the interviewed completed elementary school and 16.6% was illiterate. A 9.9% of the residents reported at least one family member having had malaria. A 98.1% of families used some method to prevent mosquito bites; those using bed nets were 94.3%. Almost 72% of families bought products for mosquito protection. A total of 537 out of 542 families agreed with the indoor residual spraying (IRS) of insecticide and a frequency of application as often as every two months was preferred.

Conclusion: Housing conditions and malaria preventive practices have improved in these rural areas in 17 years, which could be in favor of malaria elimination in this area. Information generated by this study could help in the decision making about whether to use insecticide as indoor residual spraying or to implement massive distribution of long-lasting impregnated bed nets, considering the number of bedrooms and the structure of houses in the region, which may lead to a more efficient vector control for the coastal plain of Chiapas.

Keywords: Socioeconomic factors, Attitudes, Practices, Malaria prevention, Mexico, Questionnaires, Interviews, Bed nets, Indoor residual spraying, Living conditions

* Correspondence: penilla@insp.mx

Centro Regional de Investigación en Salud Pública /Instituto Nacional de Salud Pública, Cuarta Avenida Norte y 19 Calle Poniente, C.P. 30700 Tapachula, Chiapas, México

Background

Malaria is still a disease considered endemic in 106 countries and territories, despite control efforts implemented [1,2]. Although Mexico is in the malaria pre-elimination phase [2,3], Chiapas is considered a risk area with low, seasonal and persistent transmission [4]. During 2011, 54.1% of malaria cases in Mexico were reported in Chiapas [5], and although the cases were reduced to 32.0% during 2012, Chiapas continues being the foci with the highest malaria prevalence. Sociocultural and economic factors, such as education, housing patterns and social groups, play an important role in malaria transmission [6]. Experiences with malaria have shown that prevention is better and cheaper than cure [7].

Action plans for malaria control suggest applying indoor residual spraying (IRS) of insecticide and the use of insecticide-impregnated bed nets to prevent mosquito bites, as well as monitoring transmission mainly in areas with high levels of receptivity and vulnerability [8,9]. Questionnaires have been the tool most widely used to know factors related with malaria, such as sociocultural and demographic factors, and the information they provide helps to know and evaluate malaria control measures implemented by the control programmes, as well as to provide valuable insights necessary to guiding the implementation of malaria health promotion strategies.

In the context of the possible pre-elimination of malaria in the Mesoamerican region, and considering that vector control has been relaxed during the last decade in Mexico, it is important to determine how keen the communities are in receiving and participate with the malaria control measures if they were again implemented, as well as to know the differences in the sociocultural and demographic factors in places where the control programme used to have a continuous activity.

The aim of this study was to investigate the socioeconomic factors, attitudes and practices in the prevention of vector borne diseases like malaria, by the application of a questionnaire. Additionally, information about education levels was collected in twenty villages from the coastal plain of Chiapas, México. Responses were discussed and results compared to those given 17 years ago by villagers from the same region.

Methods

Study area

This study was conducted in the coastal plain of Chiapas, Mexico, in the same area where a similar questionnaire was applied 17 years ago [10] previously to a mosaic and rotation application trial of insecticides. Twenty villages from six municipalities were selected: El Fortín, Chochohuitl, El Topón, Las Coaches and Buena Vista from municipality Pijijiapan; Barrita Pajón, 10 de Abril, El Castaño and Juan Escutia from municipality Mapastepec;

15 de Abril, Las Lauras and Río Arriba Embarcadero from municipality Acapetahua; Vicente Guerrero, Emiliano Zapata and 19 de Abril from municipality Mazatán; Lázaro Cárdenas from municipality Tapachula and; 15 de Septiembre, Conquista Campesina, Brisas del Mar and Emiliano Zapata from municipality Suchiate (Figure 1).

Data collection

The study took place in January 2012. A questionnaire (approved by the Public Health Ethics Committee of the National Institute of Public Health, México) was administered by six trained survey personnel in a face-to-face interview with an available adult member of the family (Additional file 1 shows the questionnaire administered). Interviewers were distributed homogeneously inside the villages, so household's information was representative. Each interviewer randomly selected the houses per block until completed the number of questionnaires assigned. According to previous studies in this area applying questionnaires [10], where at least 30% of the households were included, in this study 30 questionnaires per village was calculated to meet at least 30% of the households, except for small villages with less than 30 houses. The aim of the study was very carefully explained to the target population, and their consent individually obtained before questionnaire was administered.

The questionnaire comprised two parts: the first part was designed to collect socioeconomic status and education level of residents, and the second part about attitude and practices associated with malaria. The questionnaires were applied in every village; each interviewer administered five, except for the small villages where the number of houses was limited (10 de Abril, Mapastepec; Chochohuitl, Pijijiapan; El Castaño, Mapastepec; Emiliano Zapata, Suchiate and; Vicente Guerrero, Mazatán), where less than 30 questionnaires were applied. A total of 542 questionnaires were retrieved after being administered to the residents of households, one corresponding to each family.

Data were collected in Excel 14 spreadsheets where averages and percentages were calculated. Results were described and concentrated in tables where a column with data obtained from a questionnaire taken in 1995 [10] in the same study area also was included. Chi² analyses (SPSS 15.0) were performed to find out if there was association between families reporting having recent malaria and their housing conditions as their malaria preventive practices. In order to conduct the statistical analyses two indexes were created: a variable named "housing condition", in which values from 0 to 3 were assigned to every house that met the following conditions: if cement-brick walls + cement floor + cement roof were present, the value was 3; two of any (cement walls,




Figure 1 The villages selected for the study (black points), are located along the coastal plain of Chiapas, Mexico, distributed in six municipalities.

floor or roof) 2; one of any of the three 1; and none of them 0. The other index created was a variable named “preventive practices”, with three values assigned for the following conditions: if they reported having bed nets and purchased any product for mosquito bites prevention the value was 2; if just reported any of the two the value was 1; and if none 0. The variable “households reporting having malaria” also was created and 0 was given for those reporting no malaria and 1 for those reporting having had malaria at any time. Results were discussed and both housing conditions and preventive practices compared statistically by Chi² analyses against those obtained 17 years ago.

Results

Socioeconomic status and education level of residents

From the total of persons interviewed, 186 (34.3%) were males and 356 (65.7%) females. Although the average of inhabitants per house was 4.34 (with a minimum of 1 and a maximum of 30 people living in a fishermen house located at Chochohuitl), most households had 2–5 people (73.6%, $n = 399$), and only 2.3% ($n = 13$) had nine or more members, the percentage per number of people per house is described in Table 1. Children under five years were present in an important percentage of the families (36.2%, $n = 196$), most of them had only one child (24.7%, Table 1), but most of the families interviewed were formed by

Table 1 Socioeconomic and education level in the coastal plain of Chiapas between 1995 and 2012

Questions	Answers	% 1995	% 2012
Gender	Female	50.4	65.7
	Male	49.6	34.3
Total of people living in the house	4	17.9	22.5
	3	13.5	19.7
	5	20.3	18.5
	1-2	4.4	16.8
	7 and > 7	26.9	12.0
	6	17.1	10.5
People younger than 5 years old	0	0	63.8
	1	44.7	24.7
	2	41.4	9.4
	3	11.9	1.5
	>4	2.0	0.6
Bedrooms	1	69.5	40.2
	2	23.5	31.9
	3	5.0	19.6
	>4	2.0	8.3
People per room in the house	2	9.6	35.2
	1	2.4	22.5
	3	16.7	19.2
	4	23.2	10.1
	>5	45.0	12.9
Walls	Block or brick	41.9	72.3
	Palm or bamboo	20.8	15.9
	Wood	34.1	10.7
Floor	Cement	43.3	90.0
	Ground	55.8	8.8
Roof	Zinc	13.8	43.9
	Straw or Palm	51.8	42.3
	Tile	24.4	10.3
	Cement	2.6	2.6
	Others	7.2	0.9
Occupation	Housewife	49.2	65.3
	Farmer	39.2	14.4
	Fisherman	3.8	10.9
	Others	7.8	9.4
	Elementary school	13.3	23.1
Schooling	Elementary school incomplete	40.0	23.1
	Illiterate	24.1	16.6
	Middle school	3.6	14.9

Table 1 Socioeconomic and education level in the coastal plain of Chiapas between 1995 and 2012 (Continued)

Read and write	14.3	10.3
High school or higher	2.2	6.8
Middle school incomplete	2.6	5.2

persons older than 5 years old (63.8%). A majority (91.7%, $n = 497$) of households had 1–3 bedrooms, with 40.2% of the households having one bedroom, and the average number of people per bedroom was 2.7 ± 1.6 (\pm SD) (Table 1).

The physical structure of the houses consisted of walls mainly made of block or brick (72.3%, $n = 392$), palm or bamboo (15.9%, $n = 86$) and wood (10.7%, $n = 58$); the floor in most households was made of cement (90.0%, $n = 488$); while the roof was made of zinc sheet (43.9%, $n = 229$), and straw or palm (42.3%, $n = 229$) (Table 1).

Females were mostly housewives (65.3%, $n = 354$ out of the total 65.7%), while 14.4% ($n = 78$) and 10.9% ($n = 59$) out of 34.3% of the males were farmers and fishermen, respectively (Table 1). A majority of the respondents (23.1%, $n = 125$) completed elementary school (Table 1) and other 23.1% had it incomplete; 16.6% ($n = 90$) were illiterate, while just 6.8% ($n = 37$) studied high school or a higher level.

There was not a significant association between households reporting or not having recent malaria and their housing conditions according to the index which included having none, only one, any of two, or the three of the following characteristics: walls, floors and roofs made of cement or brick (Table 2).

Attitude and practices associated with malaria

All respondents had heard of malaria and responded easily to the question about whether any family member

Table 2 Household reporting having malaria according to housing conditions and preventive practices

Indexes	Reported having malaria		Total	Chi ² Sig.
	No	Yes		
Housing condition	0	6.7%	0.6%	7.2%
	1	19.5%	3.2%	22.7%
	2	61.3%	6.3%	67.7%
	3	2.4%	0.0%	2.4%
Preventive practices	0	1.7%	0.2%	1.9%
	1	25.8%	3.9%	29.7%
	2	62.5%	5.9%	68.4%

Housing conditions: 3 = cement-brick walls + cement floor + cement roof; 2 = any of cement walls, floor or roof; 1 = any of the three and; 0 = none of them.
 Preventive practices: 2 = reported having bed nets and purchased any product for mosquito bites prevention; 1 = reported any of the two and; 0 = none.
 Households reported having malaria: 1 = for those reporting having had malaria and; 0 = for those reporting no malaria.

had had malaria. Such that, 54 out of 542 (9.9%) of the residents reported at least one family member having malaria at least once in their life, and 34 out of 542 (6.3%) reported having it more than 10 year ago (Table 3). More than 98.1% of families used some method to prevent mosquito bites. Out of 542 families, 511 used bed nets (94.3%, Table 3). Ninety three percent ($n = 477$) of people who had bed nets use them all year-round and only 3.1% ($n = 16$) during the high mosquito abundance season. A total of 71.8% ($n = 389$) families bought at least one product for protection against mosquitoes. The most common product purchased was bed nets (49.4%, $n = 268$) (the other

part of families only used bed nets provided by the government at not cost), while 24.2% ($n = 131$) bought mosquito coil, 23.2% ($n = 126$) bought aerosol insecticide and just 4.0% ($n = 22$) bought mosquito repellent (Table 3). A 17.5% of the total of families ($n = 95$) bought two bed nets, 12.4% ($n = 67$) bought one, 8.5% ($n = 46$) bought three, 6.5% ($n = 35$) bought four, and 3.5% ($n = 19$) bought five or more. An average of 3.7 ± 6.0 (SD) of families bought products for mosquito bites protection during the last month before the interview. Smoke from burning cardboard is used by some families to repel mosquitos indoors. Additionally, four householders referred

Table 3 Vector control practices for preventing malaria in the coastal plain of Chiapas between 1995 and 2012

Questions	Answers	1995 %	2012 %	
Households reporting at least one in the family having malaria	Never	86.3	89.3	
	More than 10 years ago	6.0	6.3	
	6-10 years ago	3.4	1.8	
	2-5 years ago	2.4	1.7	
	1 year ago	1.6	0.2	
	Do not know	0.2	0.7	
Season usage	Have bed nets	99.0	94.3	
	All year	75.7	93.0	
	Rainy season	12.0	2.8	
	High mosq. abundance	10.6	3.1	
Use of products to prevent mosquito bites	Bought products in the last year	99.4	71.8	
	Bought bed nets in the last year	98.2	49.4	
	2	28.1	17.5	
	1	4.9	12.4	
	Number bought	3	29.1	8.5
	4	21.0	6.5	
The house was sprayed	>5	17.0	3.5	
	Mosquito coil	9.6	24.2	
	Aerosol	11.4	23.2	
	Repellent	0.8	4.0	
	>1 year ago	13.5	48.9	
	Never	56.0	37.1	
Frequency that would like an IRS	1-2 months ago	2.2	4.2	
	3-4 months ago	8.0	4.1	
	9-12 months ago	2.8	2.6	
	5-6 months ago	13.9	2.4	
	7-8 months ago	3.2	0.7	
	Every 2 months	32.7	38.6	
Frequency that would like an IRS	Every 6 months	29.3	28.0	
	Every 4 months	8.4	13.3	
	Each 3 months	22.7	12.0	
	Yearly	5.6	7.4	
	Never	-	0.7	

the use of agricultural insecticides to prevent mosquito bites; two were pyrethroid-based formulations (cypermethrin and flumethrin), one organophosphate-based (parathion) and one more an organochlorine-based formulation (lindane).

To the question about when was their houses received the last indoor residual spraying (IRS) of insecticides, 48.9% of households responded that more than a year ago (Table 3), and although a 37.1% responded that never had their houses sprayed with insecticides a total of 537 families agreed with the IRS as a strategy to protect their houses from mosquito bites in the future, and only five respondents disagreed. Three who declined argued that IRS is dangerous for children, one more complained about the smell of some insecticides, and the last person complained about removing the furniture out before the insecticide application. From those who agreed with IRS, 38.6% ($n = 209$) preferred the application as often as every two months, 28.0% ($n = 152$) every six months, 13.3% ($n = 72$) every four months, 12.0% ($n = 65$) every three months and a few of them 7.4% ($n = 40$) yearly.

There was not a significant association between households reporting or not having recent malaria and their preventive practices which included none, any or both of: having bed nets and purchased any product for mosquito bites prevention (Table 2).

Discussion and conclusion

Malaria is still one of the most important vector-borne diseases in Mexico and in the Mesoamerican region. Since 1955 when national malaria eradication programme was implemented in Mexico, malaria went through several stages all depending on the effort and amount of economic resources available for the intervention activities. In Mexico, as in many parts of the world, eradication was not achieved and a control programme was implemented instead. Nowadays, because of the control efforts but also due to some other external factors, malaria in Mexico is considered to be in the pre-elimination phase. Improvements in different aspects related with this programme are necessary for better results to finally achieve the eradication of malaria. The present study approaches topics related with socioeconomic factors, attitudes and practices in the prevention of malaria, how these factors have changed in time and how the changes may have contributed to maintain malaria transmission very low or with no transmission at all, even without any or very little malaria control activities in the coastal area of Chiapas.

It is well known that there are social determinants involved on health conditions [11], therefore some diseases may be associated with poverty and/or living conditions which in part may be exacerbated by cultural practices, although these cultural practices may also be

so influencing, that the cycle of poverty is not broken [12,13]. In the case of malaria in México, the living conditions were extremely poor when malaria was one of the first cause of mortality and morbidity, also some demographic factors were determinants for the high transmission level. By the 50's and 60's, more than 75% of total population was living in rural areas [14] where educational resources and improvements in housing conditions were much lower than in cities. In those years the percentage of illiterate people exceeded the half of the population in the Chiapas state [14], a very contrasting number compared with the results of the questionnaires applied in 1995 and 2012 in the coastal plain of Chiapas, where illiteracy decreased to 24.1% and 16.6% in the families interviewed, respectively. In the same way, poor communication with cities due to lack of roads, created a problem for awareness about malaria prevention. Coupled with this, physical structure of houses was mainly made of wood, palm or bamboo, which provided an environment more conducive to the development of malaria.

At the beginning of this century, in the same study area, Rodriguez *et al.* [10] reported 43.3% of houses with cement floor and 24.89% with cement walls. That was an important change as compared to the 1950's, however, nowadays a significantly higher percentage of houses with these characteristics (90.0% and 72.3% respectively, $P < 0.01$) were found in the same area. Changes in percentages of both structures have occurred after several implementation of government programmes, such as Solidaridad (2008), Programa Nacional de Desarrollo Social (2001–2006), Programa de Mejora de Vivienda by Comisión Nacional de la Vivienda (since 2001) and most recently the programmes Vivienda Rural by FONHAPO (Fideicomiso Fondo Nacional de Habitaciones Populares), programme of Promotora de Vivienda Chiapaneca part of the government programme “Chiapas Solidario 2007–2012”. These programmes gave successful support to improve the conditions of houses in rural areas during more than a decade ago. For example, houses having a single room decreased significantly from 69.5% to 40.2% from 1995 to 2012, while the percentage of houses with two and three rooms per house increased from 23.5% to 31.9% and from 5% to 19.6%, respectively ($P < 0.01$). This increment should be considered by authorities in the future, because a greater number of rooms involve higher expenditure for vector control programmes, specifically for the amount of insecticide and/or numbers of ITNs (Insecticide-treated nets) (although ITNs could represent a cheaper control option as will be further discussed). In fact, benefits of these improvements in living conditions together with the changes of socioeconomic factors in these localities could be in part responsible in the reduction of the malaria cases. For example, 2,750 cases were reported during 1994 and only 519 during 2011 in Chiapas

[15], that means a reduction of five times in malaria cases in this state in 17 years.

This research showed that the use of bed nets were common in the communities studied (94.3% of the households). It is well known that the use of bed nets reduces the degree of human-vector contact and malaria transmission can be significantly lowered by the use of insecticide impregnated bed nets [8,16-18], as well as could decrease malaria to pre-elimination levels by reducing mosquito human baiting rates to zero as reported for *Anopheles darlingi* after an intervention in Surinam [19]. The people interviewed have acquired this knowledge during more than a decade ago, which may have contributed to prevent the malaria transmission in the region. Official epidemiological data showed a reduction of malaria cases in Chiapas from 1995 to 2011, when people did not used to have access to impregnated bed nets as much as they do today (provided by some national programmes among them the Instituto Mexicano del Seguro Social [20]). Also, according to the 2006–2012 report from the Ministry of Health [21], 350 thousand LLNs were distributed in the malarious areas in Mexico and although no cases have been reported in the study area, some villagers reported they were provided of such bed nets from some government programme. From the study of 1995 [10], 69.3% of the families reported to use some other methods as the production of smoke from burning flammable materials, to repel mosquitos. This kind of non-commercial method is more pronounced in rural areas than urban areas [22]. Nevertheless, the high percentage of villagers using any type of bed nets may represent an advantage for a high scale ITNs implementation programme. Furthermore, it is feasible that they would be keen to pay for the ITNs, if as they responded in the questionnaires, they have purchased any product to protect themselves from mosquito bites. In this study, no direct question was asked about whether the bed nets were traditional or impregnated. However, some of the interviewed people mentioned that they were provided of ITNs by some government programme (IMSS) but they still prefer not using insecticide-impregnated bed nets because of itchy skin reactions. This could contribute to a failure if ITN mass distribution is implemented for malaria elimination in the Mesoamerican region, although no other arguments against its use were mentioned, as the listed in other studies in Peru, where interviewed people reported their privacy being affected [23], as well as a perceived lack of effectiveness due to the size of the mesh [24]. A specific study to determine whether traditional bed nets are preferred as compared to ITNs would be needed in the Mesoamerican region to assess the impact of mass distribution if it is intended to be implemented.

During 1995, according to Rodriguez *et al.* [10] more than 86% of population interviewed reported that they

had never contracted malaria, while the recently questionnaires applied showed an increment to 90.1% in the responses of households referring none of the family having malaria. This could be a good indicator of the success of the malaria control programme, but also could be a consequence of the improvement in the standard of living and socio economic level, and therefore more awareness on how to prevent malaria, including how to reduce human-vector contact. When trying to compare if the improvement of socio-economic factors could be directed linked to the referred reduction in people having malaria, a house condition and preventive practices indexes were built as reported in the methodology. These indexes were compared against reporting having malaria or not, as well as with the reported during the 1995 [10] study. No dependence of having or not malaria in the different levels of “house condition” of “preventive practices” was observed ($P > 0.05$).

In conclusion, although housing conditions, education level and malaria preventive practices have improved in these rural areas in 17 years [10], and the reports by interviewees of any member having malaria seem to be not associated with housing conditions (as measured only by walls, floors and roofs made of cement) or preventive practices (as measured only by bed nets possession and purchase of any product) an improvement of malaria control programmes needs to be implemented if the elimination of malaria is aimed in the region. Information generated by this study could help in the decision making about whether to use insecticide as indoor residual spraying or to implement massive distribution of long-lasting impregnated bed nets, considering the number of bedrooms and the structure of houses in the region, which may lead to a more efficient vector control for the coastal plain of Chiapas.

Additional file

Additional file 1: KAP questionnaire applied to householders.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MMR interviewed households, collected the data in Excel, calculated percentages, filled the tables and wrote the draft of the manuscript. RPP was involved in the study design, interviewed households, started and revised manuscript drafts and conduct the discussion. JGO was involved in the study design, interviewed households and revised manuscript drafts. ADL and FS were involved in the study design and logistic and interviewed households. JLTE was involved in the statistical analysis and revised final manuscript. ADR was involved in the study design, direction of the people, interpretation of results, revised manuscript drafts and conduct the discussion. All authors read and approved the final manuscript.

Acknowledgements

The study was supported by the WHO HQNTD1206355 project award 53270. We thank to the technicians from the Centro Regional de Investigación en Salud Pública that participated with the household interviews.

Received: 24 December 2013 Accepted: 16 April 2014
Published: 23 April 2014

References

1. Hlongwana KW, Mabaso ML, Kunene S, Govender D, Maharaj R: **Community knowledge, attitudes and practices (KAP) on malaria in Swaziland: a country earmarked for malaria elimination.** *Malar J* 2009, **8**:29.
2. WHO: *World Malaria Report 2011*. Geneva: World Health Organization; 2011.
3. Betanzos AF: **Challenges and progress in the elimination of malaria in Mexico.** *Bol Med Hosp Infant Mex* 2011, **68**:147–154.
4. Gonzalez-Ceron L, Mu J, Santillán F, Joy D, Sandoval MA, Camas G, Su X, Choy EV, Torreblanca R: **Molecular and epidemiological characterization of *Plasmodium vivax* recurrent infections in southern Mexico.** *Parasit Vectors* 2013, **18**:109.
5. Centro Nacional de Vigilancia Epidemiológica y Control de Enfermedades: *Boletín Epidemiológico 2012*. Secretaría de Salud México: Dirección general de epidemiología. <http://www.epidemiologia.salud.gob.mx/dgae/boletin/indice-2012.html>.
6. Anumudu CI, Adepoju A, Adediran M, Adeoye O, Kassim A, Oyewole I, Nwuba RI: **Malaria prevalence and treatment seeking behaviour of young Nigerian adults.** *Ann Afr Med* 2006, **5**:82–88.
7. Winch PJ, Makemba AM, Kamazina SR, Lwihula GK, Lubega P, Minjas JN, Shiff CJ: **Seasonal variation in the perceived risk of malaria, implication for the promotion of insecticide-impregnated bed nets.** *Soc Sci Med* 1994, **39**:63–75.
8. Erhun WO, Agbani EO, Adesanya SO: **Malaria prevention: Knowledge, attitude and practice in a southwestern Nigerian community.** *Afr J Biomed Res* 2005, **8**:25–29.
9. Rodríguez MH, Betanzos-Reyes AF: **Plan de mejoramiento del control de la malaria hacia su eliminación en Mesoamérica.** *Salud Pública México* 2011, **53**:333–348.
10. Rodríguez AD, Penilla RP, Rodríguez MH, Hemingway J, Betanzos AF, Hernández-Avila JE: **Knowledge and beliefs about malaria transmission and practices for vector control in Southern Mexico.** *Salud Pública México* 2003, **45**:110–116.
11. Bamba C, Gibson M, Sowden A, Wright K, Whitehead M, Petticrew M: **Tackling the wider social determinants of health and health inequalities: evidence from systematic reviews.** *J Epidemiol Community Health* 2010, **64**:284–291.
12. Institute of Medicine (US) Committee on Assessing Interactions Among Social, Behavioral, and Genetic Factors in Health: *Genes, Behavior, and the Social Environment: Moving Beyond the Nature/Nurture Debate*. Edited by Hernandez LM, Blazer DG. Washington (DC): National Academies Press (US); 2006. 5, Sex/Gender, Race/Ethnicity, and Health. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK19934/>.
13. Vorster HH: **The link between poverty and malnutrition: a South African perspective.** *Health SA Gensondheid* 2010, **15**:435.
14. Instituto Nacional de Estadística, Geografía e Informática: *Censos 1950–1990*. http://www.inegi.org.mx/prod_serv/contenidos/espanol/bvinegi/productos/integracion/pais/historicas2/cienanos/EUMCIENII.pdf.
15. Dirección General de Epidemiología, Secretaría de Salud: *Anuarios de Morbilidad*. <http://www.epidemiologia.salud.gob.mx/anuario/html/anuarios.html>.
16. Lengeler C: **Insecticide-treated bed nets and curtains for preventing malaria.** *Cochrane Database Syst Rev* 2004, **2**, CD000363.
17. Lindblade KA, Gimning JE, Kamau L, Hawley WA, Odhiambo F, Olang G, ter Kuile FO, Vulule JM, Slutsker L: **Impact of sustained use of insecticide-treated bed nets on malaria vector species distribution and culicine mosquitoes.** *J Med Entomol* 2006, **43**:428–432.
18. Hill N, Lenglet A, Arnéz AM, Carneiro I: **Plant based insect repellent and insecticide treated bed nets to protect against malaria in areas of early evening biting vectors: double blind randomised placebo controlled clinical trial in the Bolivian Amazon.** *BMJ* 2007, **335**:1023.
19. Hiwat H, Mitro S, Samjhawan A, Sardjoe P, Soekhoe T, Takken W: **Collapsing of *Anopheles darlingi* populations in Suriname after introduction of insecticide-treated nets (ITNs); malaria down to near elimination level.** *Am J Trop Med Hyg* 2012, **86**:649–655.
20. Instituto Mexicano del Seguro Social: **Portal de Compras del IMSS.** 2011. <http://compras.imss.gob.mx/?P=imsscomprofich&f=10105818&pr=>.
21. Secretaría de Salud: **Informe de Rendición de Cuentas de la APF 2006–2012.** http://portal.salud.gob.mx/contenidos/transparencia/rendicion_de_cuentas/pdf/irc_ss_3.pdf.
22. Batega DW: **Knowledge attitude and practices about malaria treatment and prevention in Uganda: A literature review.** In *Health Communication Partnership based at Johns Hopkins Bloomberg School of Public Health/Center for Communication Programs*. Baltimore and Kampala; 2004.
23. Grietens KP, Muela Ribera J, Soto V, Tenorio A, Hoibak S, Aguirre AR, Toomer E, Rodriguez H, Llanos Cuentas A, D'Alessandro U, Gamboa D, Erhart A: **Traditional nets interfere with the uptake of long-lasting insecticidal nets in the Peruvian Amazon: the relevance of net preference for achieving high coverage and use.** *PLoS One* 2013, **8**:e50294.
24. Rosas-Aguirre A, Guzmán-Guzmán M, Moreno-Gutierrez D, Rodríguez-Ferrucci H, Vargas-Pacherez D, Acuña-González Y: **Posesión, retención y uso de mosquiteros tratados con insecticidas de larga duración luego de un año de su distribución en Loreto, Perú.** *Rev Peru Med Exp Salud Publica* 2011, **28**:228–236.

doi:10.1186/1475-2875-13-157

Cite this article as: Mora-Ruiz et al.: Socioeconomic factors, attitudes and practices associated with malaria prevention in the coastal plain of Chiapas, Mexico. *Malaria Journal* 2014 **13**:157.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

