

METHODOLOGY

Open Access

Multivariable analysis of host amino acids in plasma and liver during infection of malaria parasite *Plasmodium yoelii*

Erisha Saiki^{1,2}, Kenji Nagao³, Hiroka Aonuma¹, Shinya Fukumoto², Xuenan Xuan², Makoto Bannai³ and Hirotaka Kanuka^{1,2*}

Abstract

Background: Malaria is the most significant human parasitic disease, and yet understanding of the energy metabolism of the principle pathogen, *Plasmodium falciparum*, remains to be fully elucidated. Amino acids were shown to be essential nutritional requirements since early times and much of the current knowledge of *Plasmodium* energy metabolism is based on early biochemical work, performed using basic analytical techniques, carried out almost exclusively on human plasma with considerable inter-individual variability.

Methods: In order to further characterize the fate of amino acid metabolism in malaria parasite, multivariate analysis using statistical modelling of amino acid concentrations (aminogram) of plasma and liver were determined in host infected with rodent malaria parasite, *Plasmodium yoelii*.

Results and conclusion: Comprehensive and statistical aminogram analysis revealed that *P. yoelii* infection caused drastic change of plasma and liver aminogram, and altered intra- and inter-correlation of amino acid concentration in plasma and liver. These findings of the interactions between amino acids and *Plasmodium* infection may provide insight to reveal the interaction between nutrients and parasites.

Keywords: *Plasmodium yoelii*, Infection, Nutrition, Amino acid, Aminogram, Multivariable analysis

Background

Malaria, the most significant human parasitic disease, remains a major cause of morbidity, anaemia, and mortality, in particular in developing countries. Malaria currently accounts for about two to three million deaths each year and estimates have been increasing over the last three decades [1]. It has long been acknowledged that populations residing in malaria-endemic areas generally live under conditions that lead to poor nutritional status. The groups at highest risk of the adverse effects of malaria, children and pregnant women, are also most affected by malnutrition. Infection by the human malaria parasite *Plasmodium falciparum* has severe and potentially lethal

consequences for the metabolic state of the human host [2]. In all parasitic infections, there is a significant metabolic interaction between pathogen and host as the parasite diverts nutrients towards its own growth while the host exerts to maintain homeostasis. In the case of *P. falciparum* infection, the huge metabolic demands of the rapidly proliferating parasite cells, coupled with the effects of massive erythrocyte lysis, are responsible for the pathogenesis of the disease and its clinical manifestations, which include hypoglycaemia, lactic acidosis, haemolytic anaemia, haemoglobinuria and hypoargininaemia.

Despite the clinical and economic significance of *P. falciparum*, the energy metabolism of this organism is still poorly understood. Amino acids were shown to be essential nutritional requirements of the parasite a century ago [3]. For example, the severity of malaria is related to arginine concentration in plasma and L-arginine administration to cerebral malaria patients reverts endothelial dysfunction [4,5]. Extensive genomic and biochemical

* Correspondence: kanuka@jikei.ac.jp

¹Department of Tropical Medicine, The Jikei University School of Medicine, 3-25-8, Nishi-Shinbashi, Minato-ku, Tokyo 105-8461, Japan

²National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine, Inada-cho, Obihiro, Hokkaido 080-8555, Japan

Full list of author information is available at the end of the article

evidence indicate that many parts of parasite metabolism have been radically rationalized or modified. *Plasmodium* parasites are incapable of amino acid biosynthesis, relying on haemoglobin catabolism and uptake from the extracellular space [6]. Upon invading an erythrocyte, the parasite initiates a catabolic process whereby haemoglobin from the erythrocyte cytoplasm is ingested and proteolysed into its constituent amino acids in an acidic vacuole [7]. Therefore, the growth and multiplication of parasites are dependent on the presence in the extracellular solution of a set of amino acids, for example, isoleucine or methionine [8-11], that are absent or very little from adult human haemoglobin [12]. In addition, amino acids are substrate of proteins and play a central role as intermediates in nutrition metabolism, which form complicated networks, such as lipid, nitrogen and carbohydrate [13]. Thus, a novel analytic method is required to be developed in a blanket manner to understand the interaction among parasites and host amino acid metabolisms.

This study aimed to perform multivariable analysis using statistical modelling of amino acid concentrations (aminogram) [14,15] of plasma and liver of the host that was infected by rodent malaria parasite, *Plasmodium yoelii*. Plasma aminogram analysis indicates comprehensive survey of the amino acid concentrations in plasma, which has already been used in diagnosis of cancer [16]. Due to the importance of *Plasmodium* amino acid metabolism for malaria pathogenesis and as a target for most current candidate anti-malarial pharmaceuticals, it is critical to understand the dynamics of the amino acid metabolic networks in *Plasmodium* subspecies. Alternative approaches to reconstructing these networks, such as presumption from genomic data or *in vitro* biochemistry, are currently incomplete and metabolomics technologies are beginning to enable systems-level measurements of changes in metabolic activity [17,18]. This study provides a novel view to reveal the host-parasite interaction in aspects of nutrition dynamics, and the role played by these complicated pathways in maintaining the host-parasite balance.

Methods

Ethics statement

This study was carried out in strict accordance with the recommendations in the Guide for the Laboratory Animals of the Obihiro University of Agriculture and Veterinary Medicine and The Jikei University School of Medicine. The protocol was approved by the Committee on the Animal Experiments of the Obihiro University of Agriculture and Veterinary Medicine (Permit Number: 21-41 and 21-42) and The Jikei University School of Medicine (Permit Number: 23-020). All experiments using mice were performed under anaesthesia, and all efforts were made to minimize suffering in accordance with the Guidelines for Animal Experimentation of the

Japanese Association for Laboratory Animal Science and the Fundamental Guidelines for Proper Conduct of Animal Experiment and Related Activities in Academic Research Institutions under the jurisdiction of the Ministry of Education, Culture, Sports, Science and Technology, Japan.

Animals

Six- to eight-week-old male BALB/c mice (CLEA Japan) were infected with the *P. yoelii* 17XL strain by the intraperitoneal injection of infected blood. Asexual growth was monitored by Giemsa-stained smear. The animal room was maintained at a constant temperature ($23\pm 2^\circ\text{C}$) and humidity ($55\pm 10\%$) with a 12:12 hours light-dark cycle (lights on 0800-2000). Under diethylether anaesthesia, blood and liver were collected. The liver samples were immediately frozen and stored at -80°C until analysis.

Parasite infection

Donor BALB/c mice were intraperitoneally injected with 1×10^6 erythrocytes infected with *P. yoelii* 17XL strain [19,20]. Parasitaemia was monitored by counting the number of parasite-infected erythrocytes per 1,000 erythrocytes by microscopic examination of Giemsa-stained, thin (tail) blood smears.

Amino acid analysis

Animals were bled under ether anaesthesia; blood was collected into tubes containing EDTA as anticoagulant. After centrifugation, the supernatant was used for examination of plasma. Plasma samples were mixed with two volumes of 5% (w/w) trichloroacetic acid, and centrifuged (4°C , 20 min, 8,000 g) immediately after blood collection to remove precipitated protein. To prepare deproteinized tissue extracts, tissues were homogenized by using 5% trichloroacetic acid as described previously [21]. All samples were kept at 4°C during all steps to minimize chemical reactions of thiol metabolites. The amino acid concentrations were measured by an automatic amino acid analyzer (L-8800; Hitachi, Tokyo, Japan). Briefly, amino acids, separated by cation exchange chromatography, were spectrophotometrically detected after post-column reaction with ninhydrin reagent. Hydroxyproline content was determined by the method previously reported [22]. 20 amino acids (Asn, Gln, Asp, Thr, Ser, Glu, Pro, Gly, Ala, Cys, Val, Met, Ile, Leu, Tyr, Phe, His, Lys, Trp, Arg), 11 their derivative (Hydroxyproline (Hypro), Hydroxylysine (Hylys), 3-methylhistidine (3-MeHis), 1-methylhistidine (1-MeHis), ethanolamine (EOHNH₂), beta-alanine (beta-Ala), alpha-aminobutyric acid (alpha-ABA), Phosphoethanolamine (PEA), Taurine (Tau), Ornithine (Orn), Citrulline (Cit)), urea, and ammonia (NH₃) were measured in this study.




Figure 1 (See legend on next page.)

(See figure on previous page.)

Figure 1 Plasma and liver amino acid concentrations vary in host infected with *Plasmodium yoelii*. Amino acid concentrations in plasma (A) and liver (B) were analysed on five days post-infection with *Plasmodium yoelii*. Shown are amino acids concentrations of uninfected mice (blue bars), infected mice (orange bars). Each bar represents mean \pm SD of six mice. Asterisks denote: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. BALB/c mice were infected with 1×10^6 RBCs parasitized with *P. yoelii*.

Aminogram analysis

Aminogram analysis was performed as previously described [14]. All data of plasma and liver concentrations of amino acids were analysed by using R (version 2.12.1; R foundation for statistical computing, Vienna, Austria [23]) for hierarchical cluster analysis. R was also used to construct coloured blocks in aminogram representing levels of amino acid concentrations, and to draw dendrogram using Ward's method [14]. The value for each

amino acid concentration was normalized to a z score by using the mean and standard deviations of the scores.

Spearman's rank correlation coefficient analysis

Intra- or inter-correlations among amino acid concentrations in plasma and liver were tested by non-parametric equivalent Spearman's rank correlation co-efficient analysis [24,25]. The calculations were performed with R, and




Figure 2 Aminogram in plasma and liver changes with *Plasmodium* infection. Alterations of aminogram in plasma (A) and liver (B) during *Plasmodium* infection are shown. The blue- and yellow-coloured cells represent z scores calculated for each amino acid [(observed mean - normal control mean)/normal control SD] in plasma and liver, respectively. Cluster analyses are applied to amino acids as described in the Methods. The z scores are as follows: yellow, positive changed; blue, negative changed; black, low variation. Coloured bars represent the mice condition and category of amino acids as follows; grey, uninfected mice; orange, *P. yoelii*-infected mice; light blue, essential amino acid for *P. falciparum*; pink, essential amino acid for mouse; green, non-essential amino acid for mouse; red, amino acid derivatives, urea, and ammonia.

the results of pair-wise correlations are shown as heat map ($|\rho| \geq 0.7, p < 0.05$).

Results

Plasma and liver amino acid concentrations vary in host infected with *Plasmodium yoelii*

To investigate amino acid metabolic profile in host during *Plasmodium* infection, concentrations of 20 amino acids, 11 their derivative, urea, and ammonia in plasma and liver were analysed in *P. yoelii*-infected mice whose parasitaemia was $84.5 \pm 3.3\%$. Ten plasma amino acids (Val, Leu, Tyr, Phe, EOHNH₂, His, Pro, Asp, Glu, Ala) and NH₃ significantly increased, and five amino acids (Cit, Cys, Met, 1-MeHis, Arg) significantly decreased (Figure 1A). On the other hand, 21 liver amino acids (PEA, Thr, Asn, Cit, α -ABA, Met, Ile, Leu, Tyr, Phe, β -Ala,

Trp, EOHNH₂, Lys, Arg, HyPro, Pro, Tau, Glu, Ala) and NH₃ significantly increased, and no amino acids significantly decreased (Figure 1B). Plasma and liver aminogram analysis based on heat-map [26] and cluster analysis [27] was then performed, demonstrating that individual plasma and liver aminogram were drastically altered by *P. yoelii* infection as well as forming infection-dependent cluster (Figures 2A and B). These data suggested that *Plasmodium* infection causes drastic alterations in host amino acid metabolism, potentially in a common pattern specific and characteristic of *Plasmodium* infection.

Intracorelation of aminogram variation in each tissue

To better understand the relationship between *Plasmodium* parasites and host amino acid levels, multivariate-





Figure 3 Intracorelation of aminogram variation in each tissue. (A) A schematic diagram of the experiment to investigate intracorelation of aminogram variation in plasma and liver. (B) Heat map of the correlation matrix among amino acids in plasma and liver. The plot summarizes the Spearman's rank correlation coefficients between 20 amino acids, 11 their derivative, urea, and ammonia in plasma and liver of mice. Coloured panel represents significant and strong correlation ($|\rho| \geq 0.7, p < 0.05$) between amino acid concentrations in plasma or liver of uninfected (upper) and infected (lower) mice. Each colour indicates as follows; green, positively correlated in plasma; magenta, positively correlated in liver; light blue, negatively correlated in liver; white, not correlated.

correlation analysis of amino acid concentrations in each of plasma and liver was performed. Spearman's rank correlation co-efficient analysis was applied to plasma and liver samples from mice infected with *P. yoelii*. The scatterplots of 2 plasma amino acids show various patterns, such as positive and negative linear correlations or no correlations (data not shown). The results were summarized as heat-map based on Spearman's rho and *p*-value (Figures 3A and B). In uninfected mice, the numbers of positively correlated pairs in amino acid concentrations were 67/528 (12.6%) in plasma and 31/528 (5.87%) in liver. In infected mice, the number of positively correlated pairs in plasma decreased to 28/528 (5.30%), while the number in liver increased to 81/528 (15.3%). Furthermore, Spearman's test between parasitaemia and alteration of amino acid concentrations showed positive (Asp, Gly, His) and negative correlation (Tyr) ($|\rho| \geq 0.7$, $p < 0.05$). These results indicated that *Plasmodium* infection

induced alterations in intra-correlations of amino acid concentrations in each tissue.

Inter-correlation of aminogram variation between plasma and liver

The amino acid transportations among organs are inter-mediated by plasma. In order to acquire further information about the external interaction between plasma and liver, Spearman's test for comparing correlations between plasma and liver amino acid concentrations was carried out. The number of positively correlated pairs in amino acid concentrations between plasma and liver were 16/1089 (1.47%) in uninfected mice, and 16/1089 (1.47%) in infected mice (Figures 4A and B). No pairs of negatively correlated were determined in uninfected mice, while pairs in infected mice were 23/1089 (2.11%). Interestingly, correlated pairs in uninfected mice did not match to any pairs in infected mice. These results indicated



that parasite infection drastically changed relationships between plasma and liver amino acid concentrations.

Discussion

Host-pathogen interactions in malaria rely on an exchange of nutrients, specifically amino acids, between erythrocyte and parasites. Amino acid concentrations of plasma were constantly kept in a set range to maintain amino acid homeostasis, although the results indicated that the concentrations are highly varied during *Plasmodium* infection (Figure 1A). The alterations of plasma amino acid concentrations by parasite infection are considered to be the results of export/import promotion of amino acids from/to erythrocytes, erythrocyte rupture, immunological enhancement, and acceleration of haematopoiesis. Since the changes of amino acid concentrations in liver influenced on plasma amino acid concentrations (Figure 1B), the variations of amino acid concentrations in other organs may also alter them. Furthermore, the enormous complexity of nutrition network makes it difficult to maintain amino acid homeostasis during parasite infection. Therefore, aminogram may lose the constancy among individual hosts during *Plasmodium* infection, although plasma and liver aminogram kept the potentially homogeneous pattern (Figures 2A and B). Amino acids are transported through blood, and their metabolism plays a central role in nutrient metabolisms; in other words, plasma aminogram indirectly reflects the metabolic dynamics in the host [28]. Consequently, these findings suggested specific patterns of nutrition dynamics responding to *Plasmodium* infection.

Metabolomic analysis gives us deeper understanding of *Plasmodium* biology and malaria patient disease state. Aminogram analysis revealed the variational aspects of host amino acid concentrations during *Plasmodium* infection, and the results suggested that altered aminogram develop novel relationships between amino acids. Furthermore, the results indicated that *Plasmodium* infection changed the intercorrelation of amino acids in plasma as well as in liver. Since amino acids act as growth regulation factors for parasites, it is reasonable to hypothesize that there is suitable plasma aminogram for parasite proliferation. These findings suggested that *Plasmodium* infection disrupted the original link and established a new one between amino acids (Figures 4A and B); that is, infection leads to novel amino acid dynamics. Meanwhile, *Plasmodium* parasites convert property of erythrocyte into adequate condition for themselves by exporting many proteins to plasma [29], such as degraded haemoglobin [30]. In fact, positive correlation was shown between plasma aminogram and amino acid composition of haemoglobin (Spearman's $\rho=0.46$, $p<0.05$), proposing a new model that *Plasmodium* parasites adjust amino acid dynamics of host to an appropriate state for themselves.

In summary, this study demonstrated that aminogram analysis and multivariate-correlation analysis of plasma and liver amino acids were efficient, to reveal the complicated interaction between parasites and host metabolisms. Through these analyses, a view to renew the traditional concepts of interaction between parasites and host was obtained. The expansion from metagenome to metabolome [31], such as aminogram analysis, is a promising approach to characterize life phenomenon, to find new candidate anti-malarial pharmaceuticals or preventative measures, and to give fresh insights into malaria control.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

ES and HK conceived the study and wrote the paper. ES, KN, and MB carried out the aminogram analysis. ES, HA, SF and XX contributed to the data analysis and discussion. All the authors read and approved the final manuscript. MB and HK supervised the study.

Acknowledgements

We are grateful to Motomi Torii for the *Plasmodium* strain and other laboratory members for their enthusiastic participation. We are grateful to K Bandai and Y Doi for skilful technical assistance, and R Paudel and K Miura for valuable discussions. This study was supported in part by Grants-in-Aid for Scientific Research from the Japanese Ministry of Education, Science, Sports, Culture and Technology to HK and SF, the Ohyama Health Foundation to HK, and the Funding Program for Next Generation World-Leading Researchers (NEXT Programme) to HK. HA is a research fellow of the Japan Society for the Promotion of Science.

Author details

¹Department of Tropical Medicine, The Jikei University School of Medicine, 3-25-8, Nishi-Shinbashi, Minato-ku, Tokyo 105-8461, Japan. ²National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine, Inada-cho, Obihiro, Hokkaido 080-8555, Japan. ³Institute for Innovation, Ajinomoto Co Inc, Kanagawa 210-8680, Japan.

Received: 8 December 2012 Accepted: 15 January 2013

Published: 16 January 2013

References

1. Krogstad DJ: **Malaria as a reemerging disease.** *Epidemiol Rev* 1996, **18**:77-79.
2. Planche T, Dzeing A, Ngou-Milama E, Kombila M, Stacpoole PW: **Metabolic complications of severe malaria.** *Curr Top Microbiol Immunol* 2005, **295**:105-136.
3. Barry DN: **Metabolism of babesia parasites in vitro: amino acid production by Babesia rodhaini compared to Plasmodium berghei.** *Aust J Exp Bio. Med Sci* 1982, **60**:175-180.
4. Yeo TW, Lampah DA, Gitawati R, Tjitra E, Kenangalem E, McNeil YR, Darcy CJ, Granger DL, Weinberg JB, Lopansri BK, Price RN, Duffull SB, Celemajer DS, Anstey NM: **Impaired nitric oxide bioavailability and L-arginine reversible endothelial dysfunction in adults with falciparum malaria.** *J Exp Med* 2007, **204**:2693-2704.
5. Olszewski KL, Morrissey JM, Wilinski D, Burns JM, Vaidya AB, Rabinowitz JD, Llinás M: **Host-parasite interactions revealed by Plasmodium falciparum metabolomics.** *Cell Host Microbe* 2009, **19**:191-199.
6. Gardner MJ, Hall N, Fung E, White OR, Berriman M, Hyman RW, Carlton JM-R, Pain A, Nelson KE, Bowman S, Paulsen IT, James K, Eisen JA, Rutherford KM, Salzberg SL, Craig AG, Kyes SA, Chan M-S, Nene V, Shallom SJ, Suh BB, Peterson JD, Angiuoli SV, Pertea M, Allen JE, Selengut JD, Haft D, Mather MW, Vaidya AB, Martin DM, Fairlamb AH, Fraunholz MJ, Roos DS, Ralph SA, McFadden GI, Cummings LM, Subramanian GM, Mungall C, Venter JC, Carucci DJ, Hoffman SL, Newbold C, Davis RW, Fraser CM, Barrell B: **Genome sequence of the human malaria parasite Plasmodium falciparum.** *Nature* 2002, **419**:498-511.

7. Krugliak M, Zhang J, Ginsburg H: **Intraerythrocytic *Plasmodium falciparum* utilizes only a fraction of the amino acids derived from the digestion of host cell cytosol for the biosynthesis of its proteins.** *Mol Biochem Parasitol* 2002, **119**:249–256.
8. Siddiqui WA, Schnell JV: **In vitro and in vivo studies with *Plasmodium falciparum* and *plasmodium knowlesi*.** *Proc Helm Soc Wash* 1972, **39**:204–210.
9. Divo AA, Geary TG, Davis NL, Jensen JB: **Nutritional requirements of *Plasmodium falciparum* in culture, I: exogenously supplied dialyzable components necessary for continuous growth.** *J Protozool* 1985, **32**:59–64.
10. Bertagna P, Cohen S, Geiman QM, Haworth J, Koenig E, Richards WH, Trigg PI: **Cultivation techniques for the erythrocytic stages of malaria parasites.** *Bull World Health Organ* 1972, **47**:357–373.
11. Liu J, Istvan ES, Gluzman IY, Gross J, Goldberg DE: ***Plasmodium falciparum* ensures its amino acid supply with multiple acquisition pathways and redundant proteolytic enzyme systems.** *Proc Natl Acad Sci USA* 2006, **103**:8840–8845.
12. Hill RJ, Konigsberg W, Guidotti G, Craig LC: **The structure of human hemoglobin. I: the separation of the alpha and beta chains and their amino acid composition.** *J Biol Chem* 1962, **237**:1549–1554.
13. Watkins SM, German JB: **Toward the implementation of metabolomic assessments of human health and nutrition.** *Curr Opin Biotechnol* 2002, **13**:512–516.
14. Noguchi Y, Zhang QW, Sugimoto T, Furuhashi Y, Sakai R, Mori M, Takahashi M, Kimura T: **Network analysis of plasma and tissue amino acids and the generation of an amino index for potential diagnostic use.** *Am J Clin Nutr* 2006, **83**:5135–5195.
15. Noguchi Y, Shikata N, Furuhashi Y, Kimura T, Takahashi M: **Characterization of dietary protein-dependent amino acid metabolism by linking free amino acids with transcriptional profiles through analysis of correlation.** *Physiol Genomics* 2008, **34**:315–326.
16. Miyagi Y, Higashiyama M, Gochi A, Akaike M, Ishikawa T, Miura T, Saruki N, Bando E, Kimura H, Imamura F, Moriyama M, Ikeda I, Chiba A, Oshita F, Imaizumi A, Yamamoto H, Miyano H, Horimoto K, Tochikubo O, Mitsushima T, Yamakado M, Okamoto N: **Plasma free amino acid profiling of five types of cancer patients and its application for early detection.** *PLoS One* 2011, **6**:49–53.
17. Kell DB: **Metabolomics and systems biology: making sense of the soup.** *Curr Opin Microbiol* 2004, **7**:296–307.
18. Bannister LH: **The interactions of intracellular protista and their host cells, with special reference to heterotrophic organisms.** *Proc R Soc London, Ser B* 1979, **204**:141–163.
19. Herbas MS, Ueta YY, Ichikawa C, Chiba M, Ishibashi K, Shichiri M, Fukumoto S, Yokoyama N, Takeya M, Xuan X, Arai H, Suzuki H: **Alpha-tocopherol transfer protein disruption confers resistance to malarial infection in mice.** *Malar J* 2010, **9**:101.
20. Zheng W, Wang QH, Liu YJ, Liu J, Feng H, Wu JJ, Cao YM: **Distinct host-related dendritic cell responses during the early stage of *Plasmodium yoelii* infection in susceptible and resistant mice.** *Parasite Immunol* 2010, **32**:324–334.
21. Hsu JM, Anthony WL, Rider AA: **Free amino acids in plasma and tissue of rats born to underfed dams.** *Proc Soc Exp Biol Med* 1975, **148**:1087–1089.
22. Rojkind M, Kershenovich D: **Hepatic fibrosis.** *Prog Liver Dis* 1976, **5**:294–310.
23. *The R project for statistical computing*; www.r-project.org.
24. Noguchi Y, Sakai R, Kimura T: **Metabolomics and its potential for assessment of adequacy and safety of amino acid intake.** *J Nutr* 2003, **133**:20975–21005.
25. Daily JP, Scanfeld D, Pochet N, Le Roch K, Plouffe D, Kamal M, Sarr O, Mboup S, Ndir O, Wypij D, Levasseur K, Thomas E, Tamayo P, Dong C, Zhou Y, Lander ES, Ndiaye D, Wirth D, Winzeler EA, Mesirov JP, Regev A: **Distinct physiological states of *Plasmodium falciparum* in malaria-infected patients.** *Nature* 2007, **450**:1091–1095.
26. Lee JS, Heo J, Libbrecht L, Chu IS, Kaposi-Novak P, Calvisi DF, Mikiyalyan A, Roberts LR, Demetris AJ, Sun Z, Nevens F, Roskams T, Thorgeirsson SS: **A novel prognostic subtype of human hepatocellular carcinoma derived from hepatic progenitor cells.** *Nat Med* 2006, **12**:410–416.
27. Rajara S, Oono Y: **NeatMap–non-clustering heat map alternatives in R.** *BMC Bioinforma* 2010, **11**:45.
28. Holmes E, Wilson ID, Nicholson JK: **Metabolic phenotyping in health and disease.** *Cell* 2008, **134**:714–717.
29. Russo I, Babbitt S, Muralidharan V, Butle T, Oksma A, Goldberg DE: **Plasmepsin V licenses *Plasmodium* proteins for export into the host erythrocyte.** *Nature* 2010, **463**:632–636.
30. Baumeister S, Winterberg M, Przyborski JM, Lingelbach K: **The malaria parasite *Plasmodium falciparum*: cell biological peculiarities and nutritional consequences.** *Protoplasma* 2010, **240**:3–12.
31. Kimura T, Noguchi Y, Shikata N, Takahashi M: **Plasma amino acid analysis for diagnosis and amino acid-based metabolic networks.** *Curr Opin Clin Nutr Metab Care* 2009, **12**:49–53.

doi:10.1186/1475-2875-12-19

Cite this article as: Saiki et al.: Multivariable analysis of host amino acids in plasma and liver during infection of malaria parasite *Plasmodium yoelii*. *Malaria Journal* 2013 **12**:19.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

