

POSTER PRESENTATION

Open Access

# Preliminary studies of biosynthesis of heme O and heme A in intraerythrocytic stages of *P. falciparum*

Raquel Simão-Gurge<sup>1\*</sup>, Julia Cricco<sup>2</sup>, Gerhard Wunderlich<sup>1</sup>, Emilia Kimura<sup>1</sup>, Brenda Cirulli<sup>2</sup>, Alejandro Katzin<sup>1</sup>

From Challenges in malaria research: Core science and innovation  
Oxford, UK. 22-24 September 2014

## Background

Intermediates of the isoprenoid metabolism can bind to heme groups. Heme is critical and its derivatives, heme O and heme A, are the main compounds of aerobic respiration. They are synthesized by farnesylation of heme groups by “heme O synthase” (HOS or Cox10) and “heme A synthase” (HAS or Cox15). The identification of heme O and heme A and characterization of enzymes involved in their synthesis are important for exploring the possible products biosynthesized by the parasite derived from the MEP pathway. Importantly, hemes A and O are essential molecules for numerous living organisms, since they are critical components of the mitochondrial electron transport chain, catalyzing the reduction of O<sub>2</sub> to H<sub>2</sub>O. Additionally, the heme group is an important target of drugs such as artemisinin and quinolines.

## Methods

The characterization of heme O in *P. falciparum* was done by two different methods using metabolic labeling with [<sup>3</sup>H] FPP and [<sup>14</sup>C] Glycine followed by extraction and separation through a C18 column permitting to observe heme O as well as heme B biosynthesis. The plasmoidal orthologs of HOS (PfCox10) and HAS (PfCox15) genes were identified by their homology to HOS and HAS enzymes from other organisms. We then genetically tagged both genes with GFP and HA coding sequences and visualized transfectants by fluorescence microscopy.

## Results and conclusion

We observed production of heme O and heme B in blood stage parasites. Visualization tagged PfCox10 and PfCox15 suggested that both enzymes were most probably localized in the mitochondria although PfCox10 presented an unusual extension at its N-terminus.

## Authors' details

<sup>1</sup>University of São Paulo, São Paulo, Sp, Brazil. <sup>2</sup>The National University of Rosario, Rosario, Argentina.

Published: 22 September 2014

doi:10.1186/1475-2875-13-S1-P82

Cite this article as: Simão-Gurge et al.: Preliminary studies of biosynthesis of heme O and heme A in intraerythrocytic stages of *P. falciparum*. *Malaria Journal* 2014 **13**(Suppl 1):P82.

Submit your next manuscript to BioMed Central  
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>University of São Paulo, São Paulo, Sp, Brazil  
Full list of author information is available at the end of the article