POSTER PRESENTATION

Open Access

2-*tert*-butyl-primaquine exhibit potent blood schizontocidal antimalarial activity via inhibition of heme crystallization

Nhien Nguyen Thanh Thuy^{1,2*}, Huy Nguyen Tien^{2,3}, Rahul Jain⁴, Kaeko Kamei²

From Parasite to Prevention: Advances in the understanding of malaria
Edinburgh, UK. 20-22 October 2010

Background

Primaquine (PQ) is the only 8-quinolinamine available to treat the malarial parasites in the infections caused by *Plasmodium vivax* and *P. ovale*. PQ has broad range of antimalarial activities, including efficacy as a causal prophylactic, gametocytocide, and sporontocide. These encouraging pharmacological properties make PQ an ideal drug to emulate while designing new antimalarials with improved activities ([1]). The placement of a metabolically stable *tert*-butyl group at the C-2 position of a quinoline ring in PQ (2-*tert*-Butyl-Primaquine - BPQ) results in a tremendous improvement in blood schizontocidal antimalarial activity ([1,2]). Because free heme released from hemoglobin catabolism in a malarial parasite is highly toxic, the parasite protects itself mainly by crystallization of heme into insoluble nontoxic hemozoin ([3]). In this study, we investigate the mechanism of blood schizontocidal activity of BPQ.

Results

The ability of 2-*tert*-butylprimaquine to inhibit in vitro beta-hematin formation (see Table 1), to form a

complex with heme with a stoichiometry of 1:1 (see Figure 1 and Figure 2), and to enhance heme-induced hemolysis (see Figure 3) were demonstrated.

Conclusion

The results described herein indicate that a major improvement in the blood-schizontocidal antimalarial activity of 2-*tert*-butylprimaquine might be due to a disturbance of heme catabolism pathway in the malarial parasite.

Table 1 IC50 values for inhibition of *P. falciparum* growth and heme crystallization

Drug	IC50 (μM) for inhibition of	
	<i>P. falciparum</i> D6 clone growth	BH formation
CQ	0.3	15.4
PQ	ND*	ND*
BPQ	0.1	2.9

*ND, not detected.

Figure 1 (abstract P69). Job plots of heme-CQ (closed circles), heme-PQ (closed triangles), and heme-BPQ (open circles) interaction. The total final combined concentration of heme and drug in the mixtures was constant at 10 μM in 40% aqueous DMSO. The pH and the temperature were constant at pH 7.4 and 25°C. The differential absorbance at 400 nm was recorded after incubation for 30 min. Values are the means ± standard errors of the means of three independent experiments. The results are reproducible.

¹Oxford University Clinical Research Unit, Hospital for Tropical Diseases, Ho Chi Minh City, Vietnam

Figure 2 (abstract P69). Hill plots of heme-CQ (closed circles) and heme-BPQ (open circles) association. The n values correspond to individual slopes. The n and K_a values for heme-CQ association were 1.09 and $3.24 \times 10^5 \text{ M}^{-1}$, respectively. The n and K_a values for heme-BPQ association were 1.07 and $0.61 \times 10^5 \text{ M}^{-1}$, respectively.

Acknowledgements

Our sincere thanks to all members in Structural Biotechnology Laboratory, Department of Applied Biology, Kyoto Institute of Technology who have contributed to and worked on this study.

Author details

¹Oxford University Clinical Research Unit, Hospital for Tropical Diseases, Ho Chi Minh City, Vietnam. ²Department of Applied Biology, Kyoto Institute of Technology, Kyoto 606-8585, Japan. ³Department of Immunogenetics, Institute of Tropical Medicine (NEKKEN), Nagasaki University; Nagasaki 852-8523, Japan. ⁴Department of Medicinal Chemistry, National Institute of Pharmaceutical Education and Research, Sector 67, S.A.S. Nagar, Punjab 160 062, India.

Published: 16 December 2010

References

- Jain M, Vangapandu S, Sachdeva S, Singh S, Singh PP, Jena GB, Tikoo K, Ramarao P, Kaul CL, Jain R: **Discovery of a bulky 2-tert-butyl group containing primaquine analogue that exhibits potent blood-schizontocidal antimalarial activities and complete elimination of methemoglobin toxicity.** *J Med Chem* 2004, **47**:285-287.
- Jain M, Vangapandu S, Sachdeva S, Jain R: **Synthesis and blood-schizontocidal antimalarial activities of 2-substituted/2,5-disubstituted-8-quinolinamines and some of their amino acid conjugates.** *Bioorg Med Chem* 2004, **12**:1003-1010.
- Egan TJ, Combrinck JM, Egan J, Hearne GR, Marques HM, Ntenti S, Sewell BT, Smith PJ, Taylor D, van Schalkwyk DA, Walden JC: **Fate of haem iron in the malaria parasite *Plasmodium falciparum*.** *Biochem J* 2002, **365**:343-347.

doi:10.1186/1475-2875-9-S2-P69

Cite this article as: Thanh Thuy et al.: 2-tert-butyl-primaquine exhibit potent blood schizontocidal antimalarial activity via inhibition of heme crystallization. *Malaria Journal* 2010 **9**(Suppl 2):P69.

Figure 3 (abstract P69). Effects of heme-induced hemolysis by drugs. **(A)** Suspensions of erythrocytes were incubated in the presence of heme (final concentrations ranged from 0 to 20 μM) without (open circles) or with 10 μM CQ (closed circles), 10 μM PQ (open triangles), or 10 μM BPQ (closed triangles). **(B)** Suspensions of erythrocytes were incubated in the presence of 10 μM of heme with various concentrations of the drug CQ (closed circles), PQ (open triangles), or BPQ (closed triangles). Error bars indicate standard deviations.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit