

RESEARCH

Open Access



Rise of multiple insecticide resistance in *Anopheles funestus* in Malawi: a major concern for malaria vector control

Jacob M. Riveron^{1*}, Martin Chiumia², Benjamin D. Menze^{1,3}, Kayla G. Barnes¹, Helen Irving¹, Sulaiman S. Ibrahim¹, Gareth D. Weedall¹, Themba Mzilahowa² and Charles S. Wondji¹

Abstract

Background: Deciphering the dynamics and evolution of insecticide resistance in malaria vectors is crucial for successful vector control. This study reports an increase of resistance intensity and a rise of multiple insecticide resistance in *Anopheles funestus* in Malawi leading to reduced bed net efficacy.

Methods: *Anopheles funestus* group mosquitoes were collected in southern Malawi and the species composition, *Plasmodium* infection rate, susceptibility to insecticides and molecular bases of the resistance were analysed.

Results: Mosquito collection revealed a predominance of *An. funestus* group mosquitoes with a high hybrid rate (12.2 %) suggesting extensive species hybridization. *An. funestus sensu stricto* was the main *Plasmodium* vector (4.8 % infection). Consistently high levels of resistance to pyrethroid and carbamate insecticides were recorded and had increased between 2009 and 2014. Furthermore, the 2014 collection exhibited multiple insecticide resistance, notably to DDT, contrary to 2009. Increased pyrethroid resistance correlates with reduced efficacy of bed nets (<5 % mortality by Olyset[®] net), which can compromise control efforts. This change in resistance dynamics is mirrored by prevalent resistance mechanisms, firstly with increased over-expression of key pyrethroid resistance genes (*CYP6Pa/b* and *CYP6M7*) in 2014 and secondly, detection of the A296S-RDL dieldrin resistance mutation for the first time. However, the L119F-GSTe2 and *kdr* mutations were absent.

Conclusions: Such increased resistance levels and rise of multiple resistance highlight the need to rapidly implement resistance management strategies to preserve the effectiveness of existing insecticide-based control interventions.

Keywords: Malaria, Insecticide resistance, Vector control, *Anopheles funestus*, Malawi

Background

Malaria remains a major public health burden in Africa [1], notably in Malawi, where it is highly endemic with an estimated six million annual cases [2, 3]. Current malaria control efforts in Malawi rely heavily on insecticide-based interventions such as long-lasting insecticide-treated nets (LLINs) and indoor residual spraying (IRS) [4]. However, reports of increasing resistance against the main insecticides used in public health are of concern

for the continued effectiveness of these control tools. In Malawi, the concern is greater for the increasing cases of resistance against pyrethroids (the only insecticide class used in bed nets) reported in the major malaria vector *Anopheles funestus* [5–8]. Insecticide resistance is a dynamic process and resistance pattern can change rapidly with time, as reported in other vector species such as *Anopheles gambiae* [9, 10]. However, changes due to ongoing control programmes in the profiles, intensity and underlying resistance mechanisms in Malawi remain largely uncharacterized. Understanding the dynamics and evolution of the resistance pattern in such a major malaria vector is crucial for the design and implementation of successful resistance management strategies.

*Correspondence: jacob.riveron@lstm.ac.uk

¹ Vector Biology Department, Liverpool School of Tropical Medicine, Pembroke Place, Liverpool, UK

Full list of author information is available at the end of the article

Design of effective control strategies relies also on a good knowledge of the vector population in term of species composition, vectorial capacity and behaviour. Such information remains patchy in Malawi, notably in the southern region where resistance has previously been reported [6]. *An. funestus* belongs to a group of ten to 11 species morphologically indistinguishable as adults [11]. However, the local species composition of this group, their role in malaria transmission, the hybridization between these species, and its impact on the introgression of genes of interests, such as resistance genes, remains largely uncharacterized.

To fill this knowledge gap and to facilitate the design and implementation of suitable vector control strategies, this study reports an extensive investigation of the dynamic changes in resistance profile and resistance mechanisms associated with ongoing insecticide-based control interventions in Malawi between 2009 and 2014. This study reveals an increase of resistance intensity and a rise of multiple insecticide resistance in *An. funestus* in Malawi causing a reduction in bed net efficacy.

Methods

Study area and mosquito collection

Adult *Anopheles* and *Culex* mosquitoes were collected in Chikwawa district (16°1'S; 34°47'E), southern Malawi, in January 2014. Geographical details of this location have been described previously [7]. Indoor-resting, blood-fed or gravid mosquitoes were collected between 06.00 and 12.00 h inside households using electric insect aspirators, after obtaining the consent of village chiefs and house owners. Collected mosquitoes were kept until fully gravid and induced to lay eggs in individual 1.5-ml microcentrifuge tubes, as described previously [12]. All F₀ females that laid eggs were morphologically identified as belonging to either the *An. funestus* group or the *An. gambiae* complex according to a morphological key [13]. Dead adult mosquitoes and egg batches were transported to the Liverpool School of Tropical Medicine under a DEFRA license (PATH/125/2012).

Species identification

To identify the different species within the *An. funestus* group, a cocktail PCR was performed as previously described [14] after genomic DNA extraction from whole mosquitoes using the DNeasy Blood and Tissue kit (Qiagen, Hilden, Germany). In addition, 50 females belonging to the *An. gambiae* complex were identified as previously described [15] after gDNA extraction [16]. Eggs were hatched in small paper cups and larvae transferred to plastic larvae trays, according to species, for rearing as previously described [12, 17].

Plasmodium sporozoite infection rate

The *Plasmodium* infection rate of *An. funestus* group mosquitoes (167 *An. funestus*, s.s., 91 *Anopheles rivulorum*-like and 37 *An. rivulorum* females) was determined using a TaqMan assay to detect four *Plasmodium* species: *Plasmodium falciparum*, *Plasmodium vivax*, *Plasmodium ovale*, and *Plasmodium malariae* [18, 19].

Insecticide susceptibility assays

Insecticide susceptibility bioassays were performed following WHO protocols [20] at 25 ± 2 °C and 70–80 % relative humidity. Assays were carried out with at least four replicates, with 25 individuals per tube, except for *An. gambiae* s.l. where only one assay per insecticide was carried out due to limited sample size. Susceptibility to ten insecticides belonging to the four major public health insecticide classes was tested in *An. funestus*: the pyrethroids permethrin (0.75 %), deltamethrin (0.05 %), lambda-cyhalothrin (0.05 %), and etofenprox (0.05 %); the carbamates bendiocarb (0.1 %) and propoxur (0.1 %); the organochlorines DDT (4 %) and dieldrin (4 %); and, the organophosphates malathion (5 %) and fenitrothion (1 %). Insecticide-impregnated papers were supplied by the WHO with the exception of etofenprox- and propoxur-impregnated papers, which were custom prepared following the standard WHO method. Two to five-day old F₁ male and female adults were exposed to insecticide-treated papers for 60 min and then held in bioassay tubes with access to 10 % sugar solution. The mortality rate was determined 24 h after exposure. In each bioassay, a control experiment (using papers impregnated only with insecticide carrier oil) was performed following the same procedure.

Piperonyl butoxide (PBO) synergist assays

Anopheles funestus s.s. F₁ adults were pre-exposed to PBO (4 %) impregnated paper for 1 h and thereafter immediately exposed to DDT (4 %) for another hour. Mortality was scored after 24 h and compared to the results obtained using DDT without PBO.

Resistance intensity

The LT₅₀ (time required for 50 % mortality) of the Chikwawa *An. funestus* s.s. population to permethrin (0.75 %), an insecticide commonly used in impregnated bed nets, was estimated after exposure of adult females at different independent timepoints: 30, 60, 90, 120, and 180 min (25 females per tube; 1–5 replicates per timepoint because of limited sample size). Because no insecticide-susceptible *An. funestus* s.s. strain was available, the resistance intensity was estimated by comparing the results with those for the *An. gambiae*-susceptible strain Kisumu (LT₅₀ 7.7 min) [21].

Bed net efficacy estimate using cone assays

In order to check the efficacy of conventional bed nets against the Chikwawa *An. funestus* population, a 3-min exposure bioassay was performed following WHO guidelines with minor modifications [22]. Five replicates of ten F₁ females (2–5 days old) were placed in plastic cones attached to two commercial nets: the Olyset® Net (containing 2 % permethrin) and the Olyset® Plus net (containing 2 % permethrin combined with 1 % of the synergist PBO). Three replicates of ten F₁ mosquitoes (2–5 days old) were exposed to an untreated net as a control. After 3 min exposure, the mosquitoes were placed in paper cups with cotton soaked in 10 % sugar solution. Knockdown was scored 1 h after exposure and mortality 24 h after exposure.

Gene expression profile of major *Anopheles funestus* insecticide resistance genes

Previous efforts to characterize the mechanisms of insecticide resistance in Malawian *An. funestus* s.s. populations revealed that pyrethroid resistance is mainly driven by key cytochrome P450s genes such as *CYP6P9a*, *CYP6P9b* and *CYP6M7* [7, 23]. The expression profiles of these three genes and one glutathione-S transferase gene (*GSTe2*) associated with *An. funestus* DDT resistance in West Africa [24], were assessed using quantitative reverse transcription PCR (qRT-PCR). These expression profiles were compared to those obtained from the 2009 sample [6]. Total RNA was extracted from three batches of ten F₁ permethrin or DDT-resistant females collected in 2014. RNA extraction, cDNA synthesis, qRT-PCR reactions and analysis were conducted as previously reported [7, 25]. The relative expression and fold-change of each target gene in 2014 and 2009 relative to the susceptible strain FANG was calculated according to the 2^{-ΔΔCT} method, incorporating PCR efficiency [26] after normalization with the housekeeping genes RSP7 (ribosomal protein S7; VectorBase ID: AFUN007153-RA) and actin (VectorBase ID: AFUN006819).

Investigation of the role of knockdown resistance mutations in pyrethroid and DDT resistance

A fragment spanning a portion of intron 19 and the entire exon 20 of the voltage-gated sodium channel gene (VGSC), including the 1014 codon associated with resistance in *An. gambiae*, was amplified and sequenced in ten permethrin-resistant and ten susceptible mosquitoes from Chikwawa and the same for DDT. PCR, sequencing and analysis were carried out as previously described [17, 27]. All DNA sequences were submitted to GenBank (Accession Number: KR337655:337726).

A TaqMan assay was performed to genotype the L1014F *kdr* mutation in *An. gambiae* s.l. samples according to Bass et al. [28]. Additionally, a fragment of the

VGSC gene spanning exon 20 was sequenced in ten field-collected *An. gambiae* s.l. for further confirmation.

Genotyping of L119F-GSTe2 and A296S-GABA receptor mutations

To assess the role of the L119F-GSTe2 mutation in DDT resistance in Chikwawa, a TaqMan assay was used to genotype 40 F₀ field-collected mosquitoes and 20 DDT-susceptible and 20 DDT-resistant F₁ mosquitoes, as described previously [24]. Likewise, the role in dieldrin resistance of the A296S-RDL mutation was assessed by a newly designed TaqMan assay used to genotype 40 F₀ field-collected female mosquitoes and 20 dieldrin-susceptible and 16 dieldrin-resistant F₁ mosquitoes. The primer and reporter sequences are shown in Additional file 1: Table S1.

Results

Mosquito species composition

More than 3000 blood-fed mosquitoes (90 % *An. funestus*, 5 % *An. gambiae* s.l. and 5 % *Culex* spp.) were collected in Chikwawa in January 2014. Of these, 512 gravid female adult *An. funestus*, 86 *An. gambiae* s.l. and 32 *Culex quinquefasciatus* were placed in 1.5-ml microcentrifuge tubes and forced to lay eggs (using a forced egg-laying method).

From the 512 females of the *An. funestus* group, 263 females laid eggs and 249 females did not. Of the 263 females that laid eggs, *An. funestus* s.s. represented 60.1 % followed by *An. rivulorum*-like (16.3 %), *An. rivulorum* (10.6 %) and *Anopheles parensis* (0.76 %) (Fig. 1a; Additional file 2: Figure S1). Surprisingly, 32 individuals (12.2 %) were hybrids: 16 *An. funestus* s.s./*An. rivulorum*-like (6.1 %), one *An. funestus* s.s./*An. rivulorum* (0.4 %), and 15 *An. rivulorum*/*An. rivulorum*-like (5.7 %). Analysis of a batch of females that did not lay eggs (139 of the 249 females) detected the same species but at significantly different proportions. Among these, *An. rivulorum*-like was predominant (60.8 %), followed by *An. funestus* s.s. (15.2 %), *An. rivulorum* (11.4 %), the hybrids *An. rivulorum*/*An. rivulorum*-like (10.1 %), and *An. funestus* s.s./*An. rivulorum*-like (2.5 %) (Fig. 1b). In the case of *An. gambiae* complex mosquitoes, all 50 F₀ females tested were identified as *Anopheles arabiensis*.

Plasmodium infection rate

Only *P. falciparum* parasites were detected in the mosquitoes. They were detected in 4.8 % (8/167) of *An. funestus* s.s. and 1.07 % (1/93) of *An. rivulorum*-like.

Insecticide susceptibility assays

Only *An. funestus* s.s. F₁ progeny were successfully reared, as both *An. rivulorum* and *An. rivulorum*-like did not generate enough individuals for bioassay tests.




Fig. 1 Species composition within the *Anopheles funestus* group in Chikwawa. **a** Females that laid eggs. **b** Females that did not lay eggs

Anopheles funestus s.s. showed multiple resistance against different insecticide classes for both females and males. Resistance was particularly high against all pyrethroids, with very low mortality rates for permethrin (type I; ♀ 13 ± 5.3 % mortality) and deltamethrin (type II; ♀ 1.8 ± 1.8 %) and lambda-cyhalothrin (type II; ♀ 4.5 ± 0.3 %), whereas high resistance was observed against the non-ester pyrethroid etofenprox with no mortality.

Similarly, this population is highly resistant to the carbamates bendiocarb (♀ 30.1 ± 5.1 % mortality) and propoxur (♀ 14.4 ± 7.2 %). Noticeably, the Chikwawa

population has developed resistance to organochlorines. Unlike in 2009, it is now resistant to DDT (♀ 69.9 ± 5.7 % mortality) and to dieldrin (♀ 83.9 ± 0.9 %). However, it remains fully susceptible to organophosphates, with 100 % mortality for fenitrothion and 98.3 % for malathion (Fig. 2a).

Resistance intensity

The LT_{50} of the Chikwawa population for permethrin was 143.5 min (females) (Fig. 2b) resulting in a resistance ratio of 18.6 (compared to susceptible *An. gambiae*).





Fig. 2 Resistance profile of the Chikwawa *An. funestus s.s.* population. **a** Mortality rates for different insecticides. **b** Timepoint mortality rates for permethrin. Error bars represent standard error of the mean

Change in resistance intensity between 2009 and 2014

Comparison of resistance levels between 2009 and 2014 reveals that the resistance intensity has significantly increased in the *An. funestus* population of Chikwawa, notably for pyrethroids. For example, for females, the mortality on exposure to deltamethrin has decreased by 40.5 % (from 42.3 % in 2009 to just 1.8 % in 2014) and by 34.2 % for permethrin (from 47.2 % in 2009 to 13 % in 2014). Similar decreases of mortality rates are observed for the other insecticides, including the carbamate bendiocarb (29.9 % reduction from 60 % in 2009 to 30.1 % in 2014) and to a lesser extent the organochlorine DDT, with an 18.9 % reduction (87.8 % in 2009 to 69.9 % in 2014). Surprisingly, resistance to organochlorines also now extends to dieldrin, with a 16.1 % reduction in mortality from 100 % in 2009 to 83.9 % in 2014 (Fig. 3a).

Insecticide susceptibility in *Anopheles arabiensis* and *Culex quinquefasciatus*

While the primary focus in this study is *An. funestus*, it is important to assess the impact of insecticide-based control interventions on other vector species, not least to compare differential effects of interventions among species. For this reason, this study also determined the insecticide resistance profiles of *An. arabiensis* and *C. quinquefasciatus*, two mosquito species that occur alongside *An. funestus* in Chikwawa, although in lower numbers. *Anopheles arabiensis* females were highly resistant to the pyrethroid permethrin (♀ 14.3 % mortality) and moderately resistant to the organochlorine DDT (♀ 87.5 % mortality), but fully susceptible to the carbamate bendiocarb (♀ 100 % mortality) (Fig. 3b). Significantly higher and multiple resistance to all four insecticide classes was observed in *C. quinquefasciatus* (Fig. 3c). For example, no mortality was observed for DDT.



Synergist assay for DDT resistance


To test whether DDT resistance was mediated by the activity of cytochrome P450 genes, mosquitoes were exposed to DDT following exposure to the synergist PBO, an inhibitor of the activity of P450s. The PBO synergist assay revealed a high recovery of susceptibility to DDT after pre-exposure for 1 h to PBO (♀ 95.9 ± 3 %; ♂ 97.30 ± 1.2 % mortality) suggesting that cytochrome P450 genes might be playing a important role in this resistance.

Insecticide-impregnated bed net efficacy assays

A nearly complete loss of efficacy was observed for the Olyset® Net (2 % permethrin), with only 2 % mortality after 3 min exposure. A higher but not full efficacy was observed for the Olyset® Plus (2 % permethrin plus 1 % PBO) net with 67 % mortality (Fig. 3d). These results support a key role for cytochrome P450s in the resistance to pyrethroids.

Transcriptional profiling of metabolic resistance genes

Significant over-expression was observed in Chikwawa in 2014 for the cytochrome P450s *CYP6P9a* and *CYP6P9b*, known to confer pyrethroid resistance in *An. funestus* [7] compared to the susceptible strain FANG [fold change (FC) 70.6 and FC 50.3, respectively]. Comparison with 2009 levels revealed that the expression levels of these two genes have increased by a factor of 1.45 for *CYP6P9a* and 1.57 for *CYP6P9b* although the difference was not statistically significant. However, a significant increase in the expression of another pyrethroid resistance gene, the cytochrome P450 *CYP6M7* [23], was observed in the samples collected in 2014 compared to 2009 (FC 3.62 ± 1.206, P < 0.05) (Fig. 4). Nevertheless, expression levels of *CYP6M7* remain lower in Chikwawa compared to *CYP6P9a* and *CYP6P9b*, with only a FC of 5.28 ± 1.76 in 2014. No change was observed for the glutathione-S transferase, *GSTe2* as expression remains low (FC < 4)



contrary to West Africa where it confers DDT and permethrin resistance [24].


Role of knockdown resistance in pyrethroid and DDT resistance

Neither the 1014 *kdr* mutation nor any mutation was detected from exon 20 of the VGSC gene (Additional file 3: Table S2). A clustering of haplotypes according to resistance phenotypes was observed for permethrin-exposed samples (Fig. 5) but not for the DDT-exposed samples (Additional file 4: Figure S2A, B), suggesting the presence of a novel *kdr* mutation in this population associated with permethrin resistance.

No 1014F mutation was detected in permethrin-resistant *An. arabiensis* by TaqMan or sequencing of the exon 20 of VGSC as previously reported in other populations of this species [29].

Role of the L119F-GSTe2 in DDT resistance

All 40 F₀ and 40 F₁ (20 DDT resistant and 20 susceptible) mosquitoes genotyped were homozygous for the susceptible L119 allele (codon CTT) indicating that DDT resistance in Chikwawa is not conferred by the L119F-GSTe2 mutation.



Detection of the A296S *RDL* mutation

Genotyping of 38 field-collected females detected the 296S *RDL*-resistant allele for the first time in a southern African *An. funestus* population at a frequency of 10.5 % and all occurring as heterozygotes. A significant association was observed between the 296S-resistant allele and dieldrin resistance as all susceptible mosquitoes were homozygous for the A296-susceptible allele whereas all resistant mosquitoes were heterozygous for 296S/A296 (odds ratio = infinity; $P < 0.0001$).

Discussion

Assessing the impact of ongoing insecticide-based control interventions on natural populations of malaria vectors is important for the design of suitable resistance management strategies. The detection in this study of a significant increase of resistance levels and the rise of multiple insecticide resistance over a 5-year period in southern Malawi provides important information on the dynamics and evolution of insecticide resistance in a major malaria vector in an area with ongoing vector control.

A complex vector population represents a challenge for malaria control

This study revealed that the composition of the *An. funestus* group in southern Malawi is more complex than previously reported, with four species identified and a large number of hybrids. A study in Karonga, in northern Malawi [8], found a similar diversity in the species composition, but did not detect *An. rivulorum*-like mosquitoes or hybrids [8]. The *An. funestus*-like, previously reported in Malawi [30], was not detected in this study despite the inclusion of primers to detect it in the species-typing PCR assay. Such diversity within the *An. funestus* group highlights the need for accurate species identification for this species group across Malawi to improve the reliability of entomological data generated from studies targeting *An. funestus* s.s., such as susceptibility levels to insecticides.

The high proportion of hybrids in this study suggests high levels of introgression between members of the *An. funestus* group. Such introgression could enable the exchange of genes of interest between these species, such as for susceptibility to *Plasmodium* infection or resistance to insecticides, possibly impacting upon their contribution to malaria transmission. This high hybridization rate between species of *An. funestus* group is similar to the high hybridization levels observed between *An. gambiae* s.s. and *Anopheles coluzzii* in the far west of their distribution range [31]. The underlying causes of such a high

level of hybridization in Chikwawa should be further investigated.

Plasmodium infection rate in southern Malawi

Of the species found, *An. funestus* s.s. is the most anthropophilic and endophilic mosquito [32], followed by *An. rivulorum*, which has previously been reported as a minor malaria vector in different African countries such as Tanzania [33] or Kenya [34]. In this study, the role of *An. funestus* s.s. in malaria transmission in southern Malawi is confirmed with infection rate similar to levels commonly reported for this species across Africa [35]. The role of *An. rivulorum* as a minor malaria vector could not be confirmed, but due to the low number of blood-fed females collected (only 37), its participation in malaria transmission cannot be ruled out either. Interestingly, this study shows that one *An. rivulorum*-like mosquito, considered mainly zoophilic and not involved in malaria transmission [36], was positive for *P. falciparum*. Further studies with higher numbers of samples are necessary to further assess the role of *An. rivulorum* and *An. rivulorum*-like mosquitoes in malaria transmission in southern Malawi.

Development of multiple insecticide resistance in Malawian *Anopheles funestus*

This study revealed an increase in the level of resistance in a period of 5 years and also a rise of multiple insecticide resistance in *An. funestus* s.s. in southern Malawi. This is of great concern for malaria control in this area where *An. funestus* s.s. is the predominant malaria vector. The increase in the resistance level is particularly high in the case of the main insecticides used in malaria control, such as pyrethroids and carbamates. Increased resistance intensity has also recently been reported in *An. gambiae* in Burkina Faso, where in 3 years the resistance ratio increased 1000-fold [37]. Such increase in resistance levels is a concern for the continued effectiveness of insecticide-based control interventions if suitable resistance management is not implemented. Equally, the rise of multiple insecticide resistance in the *An. funestus* population from Chikwawa is of concern as it limits the number of insecticide classes available for IRS. Indeed, the possible resistance to DDT observed in 2009 has now been confirmed in 2014, with only 69.9 % mortality. Resistance to organochlorines also now extends to dieldrin. The confirmation of DDT resistance is a sign of the evolution of resistance patterns in southern African populations of *An. funestus*. The only remaining insecticide class to which no evidence of resistance is seen in *An. funestus* is the organophosphates, which should be recommended for IRS using either malathion or pirimiphos-methyl (Actellic) [38].

Pyrethroid resistance reduces the efficacy of insecticide-treated bed nets

A worrying observation from this study is the loss of efficacy of bed nets against pyrethroid-resistant *An. funestus* populations from southern Malawi. This loss of efficacy is marked for the net treated only with permethrin (Olyset[®]) while a recovery of efficacy is observed with the net with added PBO (Olyset[®] Plus). Nevertheless, this synergist net still does not kill 35 % of mosquitoes. This loss of efficacy is similar to that observed in *An. gambiae* in Burkina Faso [37] and suggests that the effectiveness of LLINs might be compromised in areas of high pyrethroid resistance, particularly for LLINs without PBO to block the activity of resistance-associated cytochrome P450s. The results of this study recommend that in such areas of high pyrethroid resistance only nets with PBO be used.

Change in insecticide resistance mechanisms explains the evolution of resistance

The increased resistance to pyrethroids in this population was associated with increased expression of key cytochrome P450s previously shown to drive resistance, such as *CYP6P9a*, *CYP6P9b* and *CYP6M7* [7, 23]. This further supports previous observations that these genes are the main drivers of pyrethroid resistance in southern African *An. funestus*.

The rise of multiple insecticide resistance in Chikwawa was further supported by the first reported detection of the *RDL* mutation in a southern African population of *An. funestus* in association with dieldrin resistance. This mutation, highly prevalent in West Africa, was not reported in Malawi in 2009 or in any other southern African country [39]. However, the frequency of the 296S-resistant allele is still relatively low (10.52 %), suggesting recent introduction of the allele either as a result of gene flow from populations of other African origin or as a de novo mutation.

The absence of the L119F-GSTe2 DDT resistance mutation [24] in Chikwawa suggests that DDT resistance in southern Africa is driven by a different mechanism to that observed in West and Central Africa. The nearly full recovery of DDT susceptibility observed after PBO exposure suggests that cytochrome P450s are playing an important role, as observed in *An. gambiae* [40] or in *Drosophila* [41].

Conclusions

The increased resistance levels and rise of multiple resistance reported in here represents a serious challenge to current and future insecticide-based vector control interventions as they limit the choice of alternative insecticides for future interventions. This highlights the urgent

need to design and implement suitable resistance management strategies to ensure a continued effectiveness of existing insecticides.

Additional files

Additional file 1: Table S1. List of primers used for Taqman A296S assay and VGSC sequencing.

Additional file 2: Figure S1. Gel picture showing the species identification banding patterns obtained by PCR. Lanes are: 1, *An. funestus* s.s.; 2, *An. rivulorum*; 3, *An. rivulorum*-like; 4, Hybrid *An. funestus* s.s./*An. rivulorum*; 5, Hybrid *An. funestus* s.s./*An. rivulorum*-like; 6, Hybrid *An. rivulorum*/*An. rivulorum*-like; 7, *Anopheles parensis*.

Additional file 3: Table S2. Summary statistics for polymorphism at the sodium channel gene in susceptible and resistant permethrin and DDT *Anopheles funestus* in Chikwawa, Malawi.

Additional file 4: Figure S2. Correlation between haplotype distribution of VGSC gene and resistance phenotypes to DDT and permethrin. (A) Maximum likelihood tree of VGSC haplotypes for both DDT and permethrin-resistant and -susceptible *An. funestus* from Chikwawa; (B) For DDT alone. AL and DE denote mosquitoes alive or dead after insecticide exposure (i.e., resistant or susceptible). Perm denotes Permethrin.

Authors' contributions

CSW designed the research. JMR, MC, TM, and CSW carried out the sample collection; JMR, KGB, BDM, and SSI reared the mosquitoes; JMR performed the insecticide susceptibility and insecticide-impregnated bed net bioassays; HI and CSW performed the species identification, *Plasmodium* infection rate, transcription, genotyping, and sequencing analyses; GDW and TM contributed to data analysis and offered significant insights; JMR and CSW wrote the manuscript with contributions from all the authors. All authors read and approved the final manuscript.

Author details

¹ Vector Biology Department, Liverpool School of Tropical Medicine, Pembroke Place, Liverpool, UK. ² Malaria Alert Centre, College of Medicine, University of Malawi, Blantyre, Malawi. ³ Organisation de Coordination pour la lutte contre les Endémies en Afrique Centrale, PO Box 288, Yaoundé, Cameroon.

Acknowledgements

A Wellcome Trust Senior Fellowship in Biomedical Sciences (WT101893MA) to CSW supported this work. The authors are grateful to Kondwani Mzembe for field assistance and to Andreas Polder for technical assistance.

Compliance with ethical guidelines

Competing interests

The authors declare that they have no competing interests.

Received: 6 May 2015 Accepted: 28 August 2015

Published online: 15 September 2015

References

1. WHO. World malaria report 2013. Geneva: World Health Organization; 2014.
2. National Statistical Office NSO and ICF Macro. Malawi demographic and health survey 2010. Zomba, Malawi, and Calverton, Maryland, USA: NSO and ICF Macro; 2011.
3. Mzilahowa T, Hastings IM, Molyneux ME, McCall PJ. Entomological indices of malaria transmission in Chikwawa district, Southern Malawi. *Malar J*. 2012;11:380.
4. National Malaria Control Programme (NMCP) [Malawi] and ICF International. Malawi malaria indicator survey (MIS) 2012. Lilongwe, Malawi, and Calverton, Maryland, USA: NMCP and ICF International; 2012.

5. Hunt R, Edwardes M, Coetzee M. Pyrethroid resistance in southern African *Anopheles funestus* extends to Likoma Island in Lake Malawi. *Parasit Vectors*. 2010;3:122.
6. Wondji CS, Coleman M, Kleinschmidt I, Mzilahowa T, Irving H, Ndula M, et al. Impact of pyrethroid resistance on operational malaria control in Malawi. *Proc Natl Acad Sci USA*. 2012;109:19063–70.
7. Riverson JM, Irving H, Ndula M, Barnes KG, Ibrahim SS, Paine MJ, et al. Directionally selected cytochrome P450 alleles are driving the spread of pyrethroid resistance in the major malaria vector *Anopheles funestus*. *Proc Natl Acad Sci USA*. 2013;110:252–7.
8. Vezzenegho SB, Chiphwanya J, Hunt RH, Coetzee M, Bass C, Koekemoer LL. Characterization of the *Anopheles funestus* group, including *Anopheles funestus*-like, from Northern Malawi. *Trans R Soc Trop Med Hyg*. 2013;107:753–62.
9. Lwetoijera DW, Harris C, Kiware SS, Dongus S, Devine GJ, McCall PJ, et al. Increasing role of *Anopheles funestus* and *Anopheles arabiensis* in malaria transmission in the Kilombero Valley, Tanzania. *Malar J*. 2014;13:331.
10. Okorie PN, Ademowo OG, Irving H, Kelly-Hope LA, Wondji CS. Insecticide susceptibility of *Anopheles coluzzii* and *Anopheles gambiae* mosquitoes in Ibadan, Southwest Nigeria. *Med Vet Entomol*. 2015;29:44–50.
11. Coetzee M, Koekemoer LL. Molecular systematics and insecticide resistance in the major African malaria vector *Anopheles funestus*. *Annu Rev Entomol*. 2013;58:393–412.
12. Morgan JC, Irving H, Okedi LM, Steven A, Wondji CS. Pyrethroid resistance in an *Anopheles funestus* population from Uganda. *PLoS One*. 2010;5:e11872.
13. Gillies M, Coetzee M. A supplement to the Anophelinae of Africa South of the Sahara. *Publ S Afr Inst Med Res*. 1987;55:1–143.
14. Koekemoer LL, Kamau L, Hunt RH, Coetzee M. A cocktail polymerase chain reaction assay to identify members of the *Anopheles funestus* (Diptera: Culicidae) group. *Am J Trop Med Hyg*. 2002;66:804–11.
15. Santolamazza F, Mancini E, Simard F, Qi Y, Tu Z, della Torre A. Insertion polymorphisms of SINE200 retrotransposons within speciation islands of *Anopheles gambiae* molecular forms. *Malar J*. 2008;7:163.
16. Livak KJ. Organization and mapping of a sequence on the *Drosophila melanogaster* X and Y chromosomes that is transcribed during spermatogenesis. *Genetics*. 1984;107:611–34.
17. Cuamba N, Morgan JC, Irving H, Steven A, Wondji CS. High level of pyrethroid resistance in an *Anopheles funestus* population of the Chokwe District in Mozambique. *PLoS One*. 2010;5:e11010.
18. Bass C, Nikou D, Blagborough AM, Vontas J, Sinden RE, Williamson MS, et al. PCR-based detection of Plasmodium in Anopheles mosquitoes: a comparison of a new high-throughput assay with existing methods. *Malar J*. 2008;7:177.
19. Mulamba C, Irving H, Riverson JM, Mukwaya LG, Birungi J, Wondji CS. Contrasting Plasmodium infection rates and insecticide susceptibility profiles between the sympatric sibling species *Anopheles parensis* and *Anopheles funestus* s.s.: a potential challenge for malaria vector control in Uganda. *Parasit Vectors*. 2014;7:71.
20. WHO. Report of the WHO informal consultation tests procedures for insecticide resistance monitoring in malaria vectors, bio-efficacy and persistence of insecticides on treated surfaces. Geneva: World Health Organization: Parasitic Diseases and Vector Control (PVC)/Communicable Disease Control, Prevention and Eradication (CPE); 1998. p. 43.
21. Mawejje H, Wilding C, Rippon E, Hughes A, Weetman D, Donnelly M. Insecticide resistance monitoring of field-collected *Anopheles gambiae* s.l. populations from Jinja, eastern Uganda, identifies high levels of pyrethroid resistance. *Med Vet Entomol*. 2013;27:276–83.
22. WHO. Guidelines for laboratory and field-testing of long-lasting insecticidal nets. Geneva: World Health Organization; 2013.
23. Riverson JM, Ibrahim SS, Chanda E, Mzilahowa T, Cuamba N, Irving H, et al. The highly polymorphic CYP6M7 cytochrome P450 gene partners with the directionally selected CYP6P9a and CYP6P9b genes to expand the pyrethroid resistance front in the malaria vector *Anopheles funestus* in Africa. *BMC Genom*. 2014;15:817.
24. Riverson JM, Yunta C, Ibrahim SS, Djouaka R, Irving H, Menze BD, et al. A single mutation in the GSTe2 gene allows tracking of metabolically based insecticide resistance in a major malaria vector. *Genome Biol*. 2014;15:R27.
25. Kwiatkowska RM, Platt N, Poupardin R, Irving H, Dabire RK, Mitchell S, et al. Dissecting the mechanisms responsible for the multiple insecticide resistance phenotype in *Anopheles gambiae* s.s., M form, from Vallee du Kou, Burkina Faso. *Gene*. 2013;519:98–106.
26. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc*. 2008;3:1101–8.
27. Djouaka R, Irving H, Tukur Z, Wondji CS. Exploring mechanisms of multiple insecticide resistance in a population of the malaria vector *Anopheles funestus* in Benin. *PLoS One*. 2011;6:e27760.
28. Bass C, Nikou D, Donnelly MJ, Williamson MS, Ranson H, Ball A, Vontas J, Field LM. Detection of knockdown resistance (kdr) mutations in *Anopheles gambiae*: a comparison of two new high-throughput assays with existing methods. *Malar J*. 2007;6:111.
29. Witzig C, Parry M, Morgan JC, Irving H, Steven A, Cuamba N, et al. Genetic mapping identifies a major locus spanning P450 clusters associated with pyrethroid resistance in kdr-free *Anopheles arabiensis* from Chad. *Heredity* (Edinb). 2013;110:389–97.
30. Spillings BL, Brooke BD, Koekemoer LL, Chiphwanya J, Coetzee M, Hunt RH. A new species concealed by *Anopheles funestus* Giles, a major malaria vector in Africa. *Am J Trop Med Hyg*. 2009;81:510–5.
31. Oliveira E, Salgueiro P, Palsson K, Vicente JL, Arez AP, Jaenson TG, et al. High levels of hybridization between molecular forms of *Anopheles gambiae* from Guinea Bissau. *J Med Entomol*. 2008;45:1057–63.
32. Gillies M, De Meillon B. The Anophelinae of Africa South of the Sahara (Ethiopian Zoogeographical Region). Johannesburg: South African Institute of Medical Research; 1968.
33. Wilkes TJ, Matola YG, Charlwood JD. *Anopheles rivulorum*, a vector of human malaria in Africa. *Med Vet Entomol*. 1996;10:108–10.
34. Kawada H, Dida GO, Sonye G, Njenga SM, Mwandawiro C, Minakawa N. Reconsideration of *Anopheles rivulorum* as a vector of *Plasmodium falciparum* in western Kenya: some evidence from biting time, blood preference, sporozoite positive rate, and pyrethroid resistance. *Parasit Vectors*. 2012;5:230.
35. Coetzee M, Fontenille D. Advances in the study of *Anopheles funestus*, a major vector of malaria in Africa. *Insect Biochem Mol Biol*. 2004;34:599–605.
36. Kamau L, Koekemoer LL, Hunt RH, Coetzee M. *Anopheles parensis*: the main member of the *Anopheles funestus* species group found resting inside human dwellings in Mwea area of central Kenya toward the end of the rainy season. *J Am Mosq Control Assoc*. 2003;19:130–3.
37. Toe KH, Jones CM, N'Fale S, Ismail HM, Dabire RK, Ranson H. Increased pyrethroid resistance in malaria vectors and decreased bed net effectiveness, Burkina Faso. *Emerg Infect Dis*. 2014;20:1691–6.
38. Aikpon R, Sezonlin M, Tokponon F, Oke M, Oussou O, Oke-Agbo F, et al. Good performances but short lasting efficacy of Actellic 50 EC Indoor Residual Spraying (IRS) on malaria transmission in Benin, West Africa. *Parasit Vectors*. 2014;7:256.
39. Wondji CS, Dabire RK, Tukur Z, Irving H, Djouaka R, Morgan JC. Identification and distribution of a GABA receptor mutation conferring dieldrin resistance in the malaria vector *Anopheles funestus* in Africa. *Insect Biochem Mol Biol*. 2011;41:484–91.
40. Mitchell SN, Stevenson BJ, Muller P, Wilding CS, Egyir-Yawson A, Field SG, et al. Identification and validation of a gene causing cross-resistance between insecticide classes in *Anopheles gambiae* from Ghana. *Proc Natl Acad Sci USA*. 2012;109:6147–52.
41. Harrop TW, Sztal T, Lumb C, Good RT, Daborn PJ, Batterham P, et al. Evolutionary changes in gene expression, coding sequence and copy-number at the Cyp6g1 locus contribute to resistance to multiple insecticides in *Drosophila*. *PLoS One*. 2014;9:e84879.